

---

## Performance of Broiler Chickens Fed Differently Processed Red Kidney Beans Seed Meal (*Phaseolus Vulgaris*) as Poultry Feed Resource

\*Abdullahi, A. I., Gworgwor, Z.A., Yusuf, H.B., and Juji, C.A.

Department of Animal Science and Range Management, Modibbo Adama University,

Faculty of Agriculture, P.M.B 2076, Yola, Adamawa State, Nigeria.

\*Corresponding author: [jimetaaisha@yahoo.com](mailto:jimetaaisha@yahoo.com) ; +2348038389552

doi: <https://doi.org/10.37745/bjmas.2022.0280>

Published: August 21 2023

---

**Citation:** Abdullahi, A. I., Gworgwor, Z.A., Yusuf, H.B., and Juji, C.A. (2023) Performance of Broiler Chickens Fed Differently Processed Red Kidney Beans Seed Meal (*Phaseolus Vulgaris*) as Poultry Feed Resource, *British Journal of Multidisciplinary and Advanced Studies: Agriculture*, 4(4),63-76

---

**ABSTRACT:** *An 8 weeks feeding trial was conducted to evaluate the nutritive contribution of red kidney bean (*phaseolus vulgaris*) seed meal on growth performance, carcass and internal organ characteristics and economics of production of broiler chickens. Two hundred (Anak-2000) white strain day old broiler chicks were weight and assigned to five groups of forty chicks each. Each group was further divided into four groups of ten birds per replicate in completely randomized design. Five diets were formulated using differently processed Red kidney bean seed meal (*Phaseolus vulgaris*) at 0% in the control ( $T_1$ ) and 25% levels of inclusion for Raw (RRKBSM), Toasted (TRKBSM), boiled (BRKBSM) and Dehulled (DRKBSM) respectively. Feed and water were given ad-libitum daily throughout the feeding trial. The growth performance showed significant ( $P < 0.05$ ) difference for all the parameters. However, there were significance ( $P < 0.05$ ) for live weight, dressed weight, plucked weight, breast weight, back, thigh, drumstick, shank, kidney and lungs. Similarly cost/kg feed was higher in  $T_4$  (₦ 449.91) and lowest in  $T_2$  (₦ 424.20). Feed cost/kg weight gain was higher in  $T_5$  (₦ 440.93/kg) gain and lowest feed cost/kg gain was on  $T_4$  (₦ 308.14/kg) gain. From the foregoing, it implies that RKBSM is a potential feed resource and could be included in diets of broiler chickens when differently processed up to 25% inclusion level as protein source in broiler chicken diet for growth and performance.*

**KEYWORDS:** performance, red kidney bean, broiler chickens, carcass characteristics, economics.

---

### INTRODUCTION

Feed supply has remained a major constraint in livestock production due to the ever-increasing cost of conventional feed stuff occasioned by the competition between man and livestock (Amaefule *et al.*, 2004). The level of animal protein consumption in Nigeria is estimated at 8.27g per caput per day as against 35g per caput per day (Amaefule *et al.*, 2009). Consequently, the prices of the finished products (meat and egg) are not affordable hence the reduced protein intake. An average Nigerian does not consume enough protein of

animal origin that nourishes the body for tissue development, repairs and healthy living (FAO, 1992).

Efforts have been made to use other legume seeds as sources of protein such as Pigeon pea (Amaefule and Obioha, 2001) *Mucuna pruriens* seeds (Emenalom and Udedibie, 1998) and Jack bean (Esonu *et al.*, 1998) in monogastric diets with encouraging results. Therefore, a possible way of increasing the supply of animal products at a cheaper price is by reducing the cost of production through the use of cheaper locally available source of protein such as kidney bean in place of the imported and costly fish meal, industrial groundnut cake and or soya bean meal. Red kidney bean is rich in dietary fibre and low in fat (Krupa, 2008) Emiola *et al.* (2007) reported that red kidney bean (*Phaseolus vulgaris*) is considered as a potential component of diets of pigs and poultry.

The kidney bean (*Phaseolus vulgaris*) also known as common bean, haricot bean, navy bean, dry bean, snap bean, French bean and pinto bean are native of Central America. They are now being grown, in several parts of the world (Olomu, 2011). The bean is an herbaceous annual plant grown worldwide for its edible fruit, either the dry seed or the unripe fruit both of which are referred to as beans. Some are harvested immature and the whole pod eaten. The use of common beans in poultry and rat diets has not been too successful because of the depressing effects on performance except when anti-nutritional factors present in the seeds are deactivated (Olomu, 2011).

Red kidney bean contains high amounts of protein and energy. Its amino acid content is similar to that of soya bean except for a lower level of methionine. However, the inclusion of red kidney bean at higher amount (20%) in the diets of animal has been reported to have detrimental effects on the performance of chickens (Esonu, O. O. 2001) and rats (Apata and Ologhobo 1997). However, it has been established that heat treatment and other processing methods exert beneficial effects on the seeds of the grain legumes by destroying the anti-nutritional factors inherent in them (Balogun *et al.*, 2001). Some of the anti-nutritional factors are however thermostable, requiring different processing methods applied individually or in combination. For instance, the effectiveness of heat treatment in detoxifying Tannin, phytate and oxalate in Kidney beans has been found to be low (Emiola *et al.*, 2007). Tannin in animal nutrition has been reported to include intestinal damage, interfere with iron absorption. They are known to generally impair conversion efficiency in poultry with subsequent reduction in weight gain (Reed, 2001). Phytic acid inhibits the action of gastro-intestinal tyrosinase, trypsin, pepsin, lipase and amylase (Liener, 1980 and Khare, 2000). Oxalates are known to precipitate calcium in the gastro-intestinal tract as insoluble oxalates. Diets containing oxalates have been shown to cause calcium deficiency in animals (Hang, *et al.*, 2010). Therefore, to inactivate these anti-nutrient substances in legume seeds, emphasis has been on different processing methods. In view of the above, Red kidney Bean seeds have been processed either by toasting (Akanji *et al.*, 2003), Toasting and Boiling (Udedibe and Carnili 1997). Soaking (Akande and Fabiyi, 2010), germinating (Akande and Fabiyi, 2010), Chemical treatment Ologhobo *et al.* (1993), Supplementation with methionine (Adeyemo 2012), and supplementation with Enzyme (Abdullahi *et al.*, 2023). Birds fed cooked red kidney bean based diets has significantly higher daily weight gain than birds on

raw red kidney bean. Wu *et al.* (1996) asserted that moist treated red kidney bean had better feed consumption by birds. Thus, cooking tends to increase palatability and protein intake. Results obtained from the processed methods of red kidney beans (roasting, cooking, salting, and chemical treated) improved the amino acid availability. Lysine levels in processed seeds were higher than in the raw seeds (5.00%) except for the cooked (Ghoshit, 2015). These values obtained were higher than those for groundnut cake and soya beans meal (Olomu, 1995; Serres 1999) and Balloon (1980) reported a deficiency in methionine and lysine. Damang, *et al.* (2017) reported a higher feed intake of birds fed boiled, toasted and fermented kidney bean diets were accompanied by better utilization for growth as these birds had significantly ( $P < 0.01$ ) better feed-to-gain ratio than the chicks on all other treatments. Other processing methods need to be explored in addition to heat treatment in order to reduce the anti-nutritional factors to tolerable levels in diets of broiler Chickens.

Therefore, the objective of the study was to determine the effect of differently processed red kidney bean seed meal on growth performance and utilization by broiler chickens.

## **MATERIALS AND METHODS**

### **Location of Experiment**

The study was carried out at the Poultry Research Farm of the Department of Animal Science and Range Management of Modibbo Adama University, Yola, Girei Local Government Area of Adamawa State, Nigeria. Farm lies between latitude 9° and 11° N of the equator and longitude 10° and 14° E of the Greenwich meridian. Adamawa shares its boundaries with Taraba state to the South and West, Gombe state to the North-West and Borno state to the North. The state has tropical climate with distinct dry and wet seasons. It has an average minimum and maximum temperature of 18°C and 40°C and relative humidity of 20% and 80%. Adamawa state has an international boundary with the Cameroon Republic along its eastern boarder (Adebayo, 1999).

### **Source of Red Kidney Bean (*Phaseolus Vulgaris*)**

The red kidney bean seed was purchased from Mangu market in Mangu Local Government Area of Plateau State. The seed was subjected to the following processing methods before using it for the feeding trial.

### **Collection and Processing Red Kidney Bean Seeds**

#### **(i) Sun-dried (Raw seeds)**

Raw kidney bean seed was sun dried to reduce moisture content for sufficient feed production. Seeds procured for the study were all subjected to drying before application of other processing methods. Seeds was milled at a time in a 2mm particle size hammer mill and tagged as Raw Red Kidney Bean Diet (RRKB).

#### **(ii) Toasting (Dry Heat Treatment)**

Raw kidney seed was placed in an open metallic frying pan and the frying pan placed firmly on a locally fabricated tripod construction to serve as oven for heat using firewood. Toasting involved sprinkling water to facilitate deeper penetration. The seeds were

stirred constantly in the frying pan to ensure uniformity, and avoid charring for 60minutes. Toasting was continued until a pleasant aroma begins to emanate. Seeds was poured on the concrete floor to cool before crushing and then used as Toasted Red Kidney Bean Seed Meal, (TRKB).

#### (iii) Boiling in Alkaline Solution

Cold clean water was brought to boiling point (100<sup>0</sup>C) in a 200 litre capacity half cut drum; a modified method of boiling pigeon pea seeds was adopted. 100g of alkaline salt (KCO<sub>2</sub>) potassium bicarbonate was dissolved to produce an aqueous solution. Raw red kidney bean seeds was poured into the boiling water and covered. At end of 60 minutes, the content was drained off and boiled seeds were sundried for 4 days, before milling and then used to formulate Boiled Red Kidney Bean Diet (BRKB).

#### (iv) Dehauling

This method was carried out by use of mechanical dehauling machine. Dehauling machines are fabricated to meet the demand of various types of grains. Raw Red Kidney Bean Seeds was poured in a mechanical dehauller. At end of dehaulling, the resultant material was manually winnowed to remove the husk. The seeds were then be milled and used to formulate Dehauled Red Kidney Bean Seed Diet (DRKB).

#### Chemical analysis

Feed and processed red kidney bean seeds samples were analysed for proximate composition according to the standard described by AOAC (2010). The metabolizable energy was estimated using the formula as expressed by Ponzenga (1985).

**Table1: Proximate of Raw and Processed Red kidney Bean Seed.**

Nutrients (%)	Percentage Composition			
	RRKB	TRKB	BRKB	DRKB
Dry matter (DM)	96.44	95.99	95.55	96.99
Crude protein(CP)	24.45	23.87	22.56	24.03
Crude fibre (CF)	3.68	5.23	2.78	4.78
Ether Extract(EE)	3.69	2.87	3.43	3.33
Ash Total	5.34	3.21	2.68	4.78
Nitrogen-free extract(NFE)	56.28	46.81	54.10	60.96
*ME(Kcal/kg)	3796.50	3779.40	3.755.10	3755.90

\* ME(Kcal/kg)  $37 \times \text{CP}(\%) + 81 \times \text{EE}(\%) + 35.5 \times \text{NFE}(\%)$  (Ponzenga, 1985).

RAW – Raw Red Kidney Bean

BRKB – Boiled Kidney Bean.

TRKB – Toasted Red Kidney Bean

DRKB – Dehulled Red Kidney Bean

#### Experimental Diets

Five experimental broiler starter and finisher diets were formulated. The first diet labeled as Treatment one(T<sub>1</sub>) was the control devoid of red kidney bean seed meal while treatments 2, 3, 4 and 5 contained raw red kidney bean (RRKB), Toasted red kidney bean (TRKB), Boiled red kidney bean (BRKB) and Dehauled red kidney bean (DRKB) at 25% inclusion levels respectively as seen in Table 2 and 3.

**Table 2: Ingredients and Percentage Composition of differently processed Red kidney Broiler Starter (1-4 weeks)**

Ingredients	0% Control	25% RRKB	Dietary levels		
			25% TRKB	25% BRKB	25% DRKB
Maize	55.10	42.97	42.97	42.97	42.97
Ground nut cake	30.90	17.50	17.50	17.50	17.50
Maize bran.	3.10	3.81	3.81	3.81	3.81
Palm oil	0.10	1.35	1.35	1.35	1.35
Soya Bean meal	6.00	4.80	4.80	4.80	4.80
Red Kidney Bean	0.00	25.00	25.00	25.00	25.00
Fish meal	2.30	3.50	3.50	3.50	3.50
Bone meal	1.7	0.39	0.39	0.39	0.39
*vitamin mineral premix	0.30	0.20	0.20	0.20	0.20
salt	0.25	0.25	0.25	0.25	0.25
Lysine	0.15	0.12	0.12	0.12	0.12
Methionine	0.10	0.11	0.11	0.11	0.11
<b>Total</b>	<b>100</b>	<b>100</b>	<b>100</b>	<b>100</b>	<b>100</b>
<b>Determined Analysis</b>					
Crude protein%	22.05	22.01	22.01	22.01	22.01
ME(kcal/kg)	3007	3000	3000	3000	3000

\*Vitamin-mineral premix provided the following per kg: Vit A 1500 IU; Vit D<sub>3</sub> 3000 IU; Vit E 30 IU; Vit k 2.5mg; Thiamine B<sub>1</sub> 3mg; Riboflavin B<sub>2</sub> 6mg; Pyridoxine B<sub>6</sub> 4 mg; Niacine 40mg; Vit B<sub>12</sub> 0.02mg; Pantothenic acid 10mg; Folic 1mg; Biotin 0.08mg; Chloride 0.125 mg; Mn 0.0956g; Antioxidant 0.125g; Fe 0.024g; Cu 0.006g; Se 0.24; Co 0.24g

**Table: 3 Ingredients and percentage composition of differently processed Red kidney bean seed meal (5-8 weeks)**

Ingredients	Control 0%	Diet Inclusion Levels			
		RRKB 25%	TRKB 25%	BRKB 25%	DRKB 25%
Maize	56.10	41.96	41.96	41.96	41.96
Ground nut cake	25.00	18.88	18.88	18.88	18.88
Maize bran	6.10	4.63	4.63	4.63	4.63
Palm oil	2.80	2.80	2.80	2.80	2.80
Soya bean meal	6.00	3.46	3.46	3.46	3.46
Red kidney bean	0.00	25.00	25.00	25.00	25.00
Fish meal	1.50	1.98	1.98	1.98	1.98
Bone meal	1.70	0.61	0.61	0.61	0.61
Salt	0.30	0.21	0.21	0.21	0.21
*vit.min.premix	0.25	0.25	0.25	0.25	0.25
Lysine	0.15	0.12	0.12	0.12	0.12
Methionine	0.10	0.10	0.10	0.10	0.10
<b>Total</b>	<b>100</b>	<b>100</b>	<b>100</b>	<b>100</b>	<b>100</b>
<b>Determined Analysis</b>					
Crude protein%	19.04	19.01	19.01	19.01	19.01
ME(kcal/kg)	2923	2920	2920	2920	2920

\*Vitamin-mineral premix provided the following per kg: Vit A 1500 IU; Vit D<sub>3</sub> 3000 IU; Vit E 30 IU; Vit k 2.5mg; Thiamine B<sub>1</sub> 3mg; Riboflavin B<sub>2</sub> 6mg; Pyridoxine B<sub>6</sub> 4 mg; Niacine 40mg; Vit B<sub>12</sub> 0.02mg; Pantothenic acid 10mg; Folic 1mg; Biotin 0.08mg; Chloride 0.125 mg; Mn 0.0956g; Antioxidant 0.125g; Fe 0.024g; Cu 0.006g; Se 0.24; Co 0.2

## **Experimental Animal and Management**

Two hundred (200) day old (*Anak, 2000*) white strain broiler chicks were used for the study. The birds were managed in pens on a deep litter system throughout the study. Birds were weighed and randomly allotted to five dietary treatments of 40 birds per Treatment, 8 per replicate. Each treatment was replicated 4 times in a completely randomized design (CRD). Weighed quantities of feed were supplied to each group every morning. Left-over feeds were deducted from the quantity supplied the next morning to determine the feed intake. Thereafter birds were, weighed weekly. Feed and water were supplied *ad libitum* throughout the period of the experiment for (8) weeks.

## **Experimental design and statistical analysis**

The experimental design was completely randomized design. Data collected were subjected to one-way analysis of variance (ANOVA) as described by Steel and Torrie (1980). Mean separation were carried out as described by Duncan's multiple range test (Duncan 1955).

## **Carcass and internal Organs Evaluation**

Eight birds from each treatment, two from each replicate were randomly selected at the end of the experiment for carcass analysis (total of 40 chickens), allowed to starve overnight to empty the crop and water offered. The following morning, selected birds were weighed to obtain the live body weight. Slaughtered birds were allowed to bleed completely, de-feathered, and eviscerated to determine the carcass weight, internal organs weight which was expressed as a percentage of the live weight.

## **RESULTS**

### **Broiler starter phase (0-4 weeks)**

The response of broiler starter chicks to differently processed dietary levels of kidney bean seed meal was presented in table 4. The chicks fed diets containing BRKB had significantly ( $P<0.05$ ) highest average daily feed intake (67.23 g/bird/day) than 58.04, 53.93, 51.07 and 46.43 g for birds fed Raw, Toasted, dehaulled and Controlled diets respectively. Birds fed toasted kidney beans diets also had a significantly ( $P<0.05$ ) Higher feed intake than those fed the control diets. Chicks fed dehaulled kidney bean diet had the least feed intake. However, there was a highly significant ( $P<0.05$ ) difference for final weight (346.50, 442.25, 455.50, 456.60 and 678.25 g/bird). The BRKBMS ( $T_4$ ) had the highest (678.25 g/bird) While DRKBSM ( $T_5$ ) had the least weight. Feed conversion Ratio (FCR) did not differ significantly ( $P>0.05$ ).

The total body weight gain of chicks fed Raw, Toasted, Boiled and Dehaulled Kidney bean was significant ( $P<0.05$ ) among treatment groups. Boiled kidney bean had the highest body weight (639.75 g) and Dehaulled kidney bean (309.25 g) had the least.

**Table 4: Effects of differently processed RKBSM on performance of Broiler Chickens Starter Phase (1-4 weeks)**

Parameter	Control	RRKB(25%)	Level of Differently Processed			SEM
			TRKB(25%)	BRKB(25%)	DRKB(25%)	
	T <sub>1</sub>	T <sub>2</sub>	T <sub>3</sub>	T <sub>4</sub>	T <sub>5</sub>	
Initial Weight (g)	37.75	39.00	38.75	38.5	37.25	1.81 <sup>ns</sup>
Final Weight (g)	455.50 <sup>b</sup>	456.60 <sup>b</sup>	442.25 <sup>b</sup>	678.25 <sup>a</sup>	346.50 <sup>b</sup>	53.83 **
Total weight (g)	417.75 <sup>b</sup>	417.50 <sup>b</sup>	403.50 <sup>b</sup>	639.75 <sup>b</sup>	309.25 <sup>b</sup>	53.01 **
Total feed intake(g)	1510.00 <sup>b</sup>	1430.00 <sup>b</sup>	1625.00 <sup>ab</sup>	1882.50 <sup>a</sup>	1400.00	112.9 3*
Average daily feed intake(g)	53.93 <sup>b</sup>	51.07 <sup>b</sup>	58.04 <sup>ab</sup>	67.23 <sup>a</sup>	46.43 <sup>b</sup>	4.03*
Feed Conversion Ratio	3.63	3.4	4.0	2.9	4.5	0.30 <sup>ns</sup>

*a, b, c, d: means within the same row having different superscript are significantly different*

#### Broiler Finisher Phase (5-8 weeks)

The response of finisher broiler chickens to differently processed kidney beans studied was presented in table 5. Total feed intake of Chickens fed BKBSM was highest (3862.50 g) ( $P>0.05$ ) and lowest (3622.50 g) for DKBSM. The Average daily feed intake was also highest on BKBSM (137.70 g/bird) as compared to TKBSM (129.73 g), RKBSM (132.10 g), and DKBSM (130.80 g) which was not significant ( $P>0.05$ ). Daily weight gain was however significant ( $P<0.05$ ) with 52.29 g/bird/day for birds on BKBSM. This was followed by TKBSM (44.38 g/bird/day), Compared with (42.54 and 34.68 g/bird/day) for RRKBSM and DRKBSM. This result was observed to be different for the control (47.84 g/bird/day). FCR of 2.60 was observed for birds on BRKBSM, and 2.90 for TRKBSM as against RRKBSM and DRKBSM (3.00 and 3.70) respectively.

**Table 5: Effects of differently processed RKBSM on performance of Broiler Chickens Finisher Phase (5-8 weeks)**

Parameter	Control 0%	Levels of differently processed				SEM.
		RRKB 25%	TRKB 25%	BRKB 25%	DRKB 25%	
	T <sub>1</sub>	T <sub>2</sub>	T <sub>3</sub>	T <sub>4</sub>	T <sub>5</sub>	
Initial Weight (g)	455.50 <sup>b</sup>	456.60 <sup>b</sup>	442.75 <sup>b</sup>	678.25 <sup>a</sup>	346.50 <sup>b</sup>	53.83 <sup>**</sup>
Final Weight (g)	1795.00 <sup>b</sup>	1647.50 <sup>b</sup>	1655.00 <sup>b</sup>	2142.50 <sup>a</sup>	1317.50 <sup>c</sup>	106.59 <sup>**</sup>
Total weight (g)	1339.50 <sup>a</sup>	1191.00 <sup>ab</sup>	1242.75 <sup>ab</sup>	1464.25 <sup>a</sup>	971.00 <sup>b</sup>	97.81 <sup>*</sup>
Average daily weight gain(g)	47.84 <sup>a</sup>	42.54 <sup>ab</sup>	44.38 <sup>ab</sup>	52.29 <sup>ab</sup>	34.68 <sup>b</sup>	3.49 <sup>*</sup>
Total feed intake(g)	3854.50	3632.50	3650.00	3862.50	3622.50	208.65 <sup>ns</sup>
Average daily feed intake(g)	137.70	129.73	130.40	137.90	129.40	7.45 <sup>ns</sup>
Feed Conversion Ratio	2.70 <sup>b</sup>	3.0 <sup>ab</sup>	2.90 <sup>ab</sup>	2.60 <sup>a</sup>	3.70 <sup>a</sup>	0.21 <sup>*</sup>

*a, b, c, d: means within the same row having different superscript are significantly different*

Record on carcass yield and internal organ characteristics is presented in table: 5. The results obtained showed that there were no significant ( $P>0.05$ ) difference among treatment groups for the dressing percentage with range value 65.49, 65.55, 60.88, 71.76 and 58.9 g respectively for Control (0%), RRKB, TRKB, BRKB and DRKB. While the result for the live weight, dressed weight, plucked weight, breast weight, back, thigh, drumstick, wings, shank, kidney and lungs all show significance ( $P<0.01$ ) and ( $P<0.05$ ) differences across treatment groups.

**Table 6: Carcass yield and internal organ characteristics of broilers fed differently processed RKBSM**

Parameter	Control T <sub>1</sub> (0%)	Dietary Levels				SEM
		RRKB T <sub>2</sub>	TRKB T <sub>3</sub>	BRKB T <sub>4</sub>	DRKB T <sub>5</sub>	
Live wt (g)	1695.00 <sup>b</sup>	1547.50 <sup>b</sup>	1585.00 <sup>b</sup>	2042.50 <sup>a</sup>	1217.50 <sup>c</sup>	106.59 <sup>**</sup>
Dressed wt (g)	1107.50 <sup>b</sup>	1020.00 <sup>bc</sup>	972.50 <sup>bc</sup>	1467.50 <sup>a</sup>	735.00 <sup>c</sup>	111.39 <sup>**</sup>
Dressing %	65.49	65.55	60.88	71.76	58.91	4.40 <sup>ns</sup>
Plucked wt	1620.00 <sup>ab</sup>	1462.50 <sup>b</sup>	1365.00 <sup>b</sup>	1820.00 <sup>a</sup>	751.25 <sup>c</sup>	95.89 <sup>**</sup>
Breast wt (g)	311.50 <sup>ab</sup>	290.25 <sup>ab</sup>	299.75 <sup>ab</sup>	380.00 <sup>a</sup>	205.75 <sup>b</sup>	32.96 <sup>**</sup>
Back (g)	148.50 <sup>ab</sup>	125.00 <sup>bc</sup>	152.25 <sup>ab</sup>	167.50 <sup>a</sup>	100.75 <sup>c</sup>	11.38 <sup>**</sup>
Thigh (g)	211.00 <sup>a</sup>	161.50 <sup>b</sup>	155.00 <sup>b</sup>	207.50 <sup>a</sup>	124.50 <sup>b</sup>	14.91 <sup>**</sup>
Drumstick (g)	165.00 <sup>a</sup>	147.75 <sup>ab</sup>	141.75 <sup>ab</sup>	179.50 <sup>a</sup>	111.75 <sup>b</sup>	13.61 <sup>*</sup>
Wings (g)	134.50 <sup>ab</sup>	114.25 <sup>bc</sup>	109.50 <sup>bc</sup>	152.50 <sup>a</sup>	94.75 <sup>c</sup>	9.9 <sup>**</sup>
Shank (g)	75.75 <sup>a</sup>	75.75 <sup>a</sup>	62.75 <sup>ab</sup>	74.75 <sup>a</sup>	52.75 <sup>b</sup>	4.80 <sup>*</sup>
Crop (g)	9.00	8.25	10.50	10.50	13.50	2.7 <sup>ns</sup>
Thorax (g)	73.00	65.75	71.50	85.00	62.50	8.57 <sup>ns</sup>



Neck (g)	82.00	73.25	66.75	96.00	55.25	9.36 <sup>ns</sup>
Head (g)	52.50	46.00	36.50	51.50	36.75	5.03 <sup>ns</sup>
Pancreas	5.00	4.00	5.50	6.00	3.00	0.95 <sup>ns</sup>
Large/small intestines(cm)	135.00	112.00	128.20	119.75	90.00	12.62 <sup>ns</sup>
Gizzard wt(g)	2.20	2.61	2.80	1.95	2.29	0.22 <sup>ns</sup>
Liver (g)	2.14	2.36	2.43	2.03	2.54	0.24 <sup>ns</sup>
Kidney (g)	0.32 <sup>ab</sup>	0.84 <sup>a</sup>	0.55 <sup>b</sup>	0.39 <sup>b</sup>	0.82 <sup>b</sup>	0.11 <sup>**</sup>
Spleen (g)	0.09	0.18	0.13	0.09	0.18	0.01 <sup>ns</sup>
Lungs (g)	0.53 <sup>a</sup>	0.75 <sup>a</sup>	0.54 <sup>a</sup>	0.48 <sup>a</sup>	0.88 <sup>b</sup>	0.05 <sup>**</sup>
Proventriculus	0.63	0.83	0.70	0.65	0.90	0.089 <sup>ns</sup>
Abdominal fat	0.86	2.01	0.86	1.75	0.79	0.47 <sup>ns</sup>

*a, b, c : means within the same row having different superscript are significantly different*

**Table 7: Economics of Production in Broiler Chickens Fed Differently Processed RKBSM**

Parameter	Dietary Inclusion Levels					SEM
	Control T1(0%)	RRKB T2(25)	TRKB T3(25%)	BRKB T4(25%)	DRKB T5(25%)	
Total feed intake (bird/kg)	3.87	3.63	3.65	3.85	3.66	0.01
Cost of feed (N/kg)	110.4	116.86	116.86	116.86	116.86	0.48
Cost of feed intake (N/kg)	427.24	424.20	426.53	449.91	427.70	1.81
Total weight gain (Kg)	1.34	1.21	1.24	1.46	0.97	0.05
Feed cost/kg weight gain	318.84	350.58	343.98	308.15	440.93	1.40

## DISCUSSION

The inclusion of 25% Kidney bean seed in the diet of Chicken did affect growth in the starter phase ( $P < 0.05$ ) and the Finisher phase as reflected in Table 3 and 4. A reduced efficiency of feed utilization was observed which was also reflected in reduced weight gain in RRKB, TRKB and DRKB except for BRKB recorded a slightly higher gain weight. This agrees with report of (Wu *et al.*, 1996) Stating that boiling tends to increase palatability while increasing protein intake. It also supports earlier findings of (Ologhobo *et al.*, 1993) that nutritive value of legumes is enhanced by processing methods. The average daily feed intake in the starter phase was depressed in birds fed TRKB and DRKB. This supports the reports of (Udedibe and Carlini, 2000; Ologhobo *et al.*, 2003). Growth rate was depressed in birds fed (RRKB and DRKB) respectively. While there was improvement in average daily gain in birds fed BRKB. The depressed growth rate observed could be due to residual anti-nutritional factors in the meal. RRKB contains Trysin inhibitors, non-starch polysaccharides (NSP), Tannins, phytate and oxalate (Afolabi *et al.*, 1985; Udedibe and Carnili, 2000). The feed conversion ratio (FCR) was significantly improved in birds fed BRKB but was considerably reduced in birds fed (control), RRKB, TRKB, and DRKB diets. Improvement in (FCR) in BRKB is consistent with the findings of Ologhobo *et al.* (1993) in lima bean and Udedibe and Carnili (2000) in Jack bean. It could also be attributed to the boiling method that

destroyed all the heat labile anti-Nutritive factors and caused reduction in others and consequently an improvement in nutritive value of the diet.

The feed intake of broiler BRKBSM diet was significantly ( $P<0.05$ ) higher than those of other treatments. This corroborates an earlier postulation (Balogun, 2001) that heat treatment of dietary legume seed significantly improved palatability in feeds. Boiling had a more positive effect on the utilization of dietary kidney beans than Toasting by significantly improving feed intake and body weight gain. This is likely to be due to the deeper penetration of heat into the seeds by boiling water, whereas during toasting, the heat is usually more superficial. This agrees with the observation of Abeke (1997) that toasted beans may not be uniformly cooked as heat penetration into the seeds is not as effective as the case with boiling.

The lower plucked weight, dressed weight and dressing percentage for birds fed RRKBSM, TRKBSM, and DRKBSM diets resulted from their smaller live weight. Changes in the pancreatic size of birds fed RRKBSM and DRKBSM are in keeping with previous reports by Meyer *et al.* (1992) who observed pancreatic Enzyme activities in pigs fed diets containing kidney beans. These authors suggested that such effects are the consequences of poor protein digestibility and this interferes with systematic protein utilization resulting in insufficient amino acids for protein synthesis.

The economic analysis of feed cost (₦/kg) as well as feed cost/kg weight gain was reduced in T<sub>1</sub> (0%) diet while in T<sub>2</sub>, T<sub>3</sub>, T<sub>4</sub> and T<sub>5</sub> (25%) the cost increased. This result agrees with the findings of Najime (2003). However, this does not agree with the report of Ani and Okorie (2005) and Bawa (2003) who reported reduction in feed cost and total cost when unconventional legume seed meals were utilized in poultry diets. This is because these legumes are cheap and available without much competition. The cost of feed/kg recorded here is higher than values reported by Omage *et al.* (2006), and Sonaiya *et al.* (1986). The reason for the high cost may be attributed to the prevailing prices of major feedstuff such as Maize, Groundnut cake and Fish meal at time of carrying out the research.

## CONCLUSION

The results can be concluded that processing methods can reduce anti nutrients such as haemagglutins, tannins, saponins, oxalates can be reduced to a reliable level. Among the processing methods, boiled red kidney bean seed meal had the best positive performance which was used as a non-conventional protein source without any effect on performance, carcass and internal organ characteristics indicating that it can compare favorably with conventional plant protein sources.

## CONFLICT OF INTEREST

The authors declare no conflict of interest that may affect the outcome of the study in any way.

## ACKNOWLEDGEMENT

The authors appreciate the Modibbo Adama University Yola for the opportunity toward sourcing the Institution Based Research (IBAR) from tertiary education fund (TETFUND). We also appreciate TETFUND and the Centre for Research and Development, Modibbo Adama University, Yola.

## REFERENCES

- Abdullahi, A. I., Gworgwor, Z.A., Yusuf, H.B., and Juji, C.A. (2023): Evaluation of processed red kidney bean (*Phaseolus vulgaris L.*) meal supplemented with methionine and enzyme in diet of broiler chickens. Unpublished Ph.D. thesis, Department of Animal Science and Range Management, Faculty of Agriculture, Modibbo Adama University, Yola, Adamawa State, Nigeria.
- Abeke, G. O. (1997): Response of laying hen to dietary levels of sheep manure. Unpublished M.Sc thesis, Department of Animal Science, Ahmadu Bello University Zaria, Nigeria. Pp 55-60
- A, F. O., (1997): Response of laying hen to dietary levels of sheep manure. Unpublished M.Sc thesis, Department of Animal Science, Ahmadu Bello University Zaria, Nigeria. Pp 55-60
- Adebayo, A.A. (1999). Climate II In: Adebayo, A.A. and Tukur, A. I. (Eds). Adamawa state Paraclete Publishers Yola.
- Adeyemo, G. O (2012) Performance of broiler starters fed varying levels of dietary methionine *International Journal of Agricultural Science*, Vol.2 (2):143-148.
- Afolabi O. A., Oshuntogun, B. A., Adewusi, S. R., Fopowo, O. O., A Yorinde, F. O., Grisson, F. E. and Oke, O. L. (1985): Preliminary nutritional and chemical evaluation of raw seeds from *Mucuna Solanei*: an under-utilized food source. *Journal of Agricultural and Food Chemistry* 33, 122 – 124.
- Akande K. E., and Fabiyi E. F., (2010): Effect of Processing methods on some anti-nutritional factors in legume seeds for poultry feeding. *International Journal of Poultry Science* (10): 966 – 1001.
- Akanji, A. M., Ologhobo, A. D., Emiola, I. A. Adedeji, T. A. and Adedeji, O. S. (2003). The effects of processing Haemagglutinin and other anti-nutritional factors on jack bean (*Canavalia ensiformis L.*). Proceedings 28<sup>th</sup> Annual Conference Nigerian Society of Animal Production Ibadan, Nigeria. Pp. 189-193.
- Amaefule, K. U and F.C. Obioha (2001). Performance and nutrient utilization of broiler fed diets containing raw, boiled or dehulled pigeon pea (*Cajanus cajan*) seeds. *Nigerian Journal of Animal Production* 28(1): 31-39.
- Amaefule, K. U. C. C., Nwaokoro, and F. C. Iheukwumere, (2004): Effects of Feeding Graded Levels of Raw Pigeon Pea Seed (*Cajanus cajan*) Meal on the Performance Nutrient Retention and Carcass Characteristics of Weaner Rabbits. *Nigerian Journal of Animal Production.*, 31:194-199.
- Amaefule, K. U., Oke, U. K. and Obioha, F.C (2009): Pigeon pea (*Cajanus cajan*) seed meal in layer diet 2. Laying performance and egg quality characteristics of pullet fed raw or processed pigeon pea seed meal diet during grower and layer stage of life. *International Journal of poultry Science*, 6:445-451.

- Ani, A. O and Okorie, A. U. (2005). The effect of graded levels of dehulled and cooked castor oil bean (*Ricinus communis, L.*) meal on performance of broiler starters. *Nigerian Journal of Animal Production*. 35(1)54-60.
- AOAC (2010). Association of Analytical Chemists. Official Methods of Analysis 18<sup>th</sup> edition Washington DC.
- Apata, D. F. and A. D. Ologhobo, (1997): Trypsin inhibitor and other anti-nutritional factors in tropical legume seeds. *Tropical Science* 37:52-59.
- Balloon, S. L. (1980).Soya bean meal in poultry nutrition. America soya bean Association St. Louis Missouri, U.S.A.
- Balogun, F. T., Kaankuka, F. G. and Bawa, G. S. (2001). Effects of boiling full-fat soya bean on its amino acid profile and on performance of pigs. *Nigerian Journal of Animal Production*. 28(1): 45-51.
- Bamgbose, A. M., Nwokoro, S. O. and O. O. Tewe, (2003): Serum Metabolites and Nutrients Utilization of Broilers Fed Extruded Full Fat Soyabean Diets. *Journal of Arid Agriculture*.13:167 – 171.
- Bawa, G. S. (2003). Effects of cooking time on the levels of anti-nutritional factors and nutritive value of dolichos lablab (*Lablab purpureusC. V. rongai*) seeds for pigs. Unpublished Ph.D. Dissertation, Dept. of Ani. Sci. ABU, Zaria.
- Damang, P. J. Tuleun, C. O. Oluremi, O. I. A. and Carew, S. N. (2017). Performance of broiler chickens fed diets containing kidney bean (*Phaseolus vulgaris L.*) seeds subjected to various processing methods.
- Duncan, D. E. (1955): Multiple range and multiple F-tests. *Biometrics*. 11:1-42.
- Emenalom, O. O., and A. B. I. Udedibie (1998).Effect of dietary raw, cooked and toasted *Mucuna pruriens* seeds (velvet bean) on the performance of finisher broilers. *Nigerian Journal of Animal Production*. 25:115-118.
- Emiola, I. A. and Ologboho, A. D. (2007). Nutritional assessment of raw and differently processed underutilized legume seeds in broiler diets. *Journal of Animal and Veterinary Advance*, 5(2):96-105.
- Esonu, B. O., Emenalom, O. O., Udedibie, A. B. I., Herbert, U., Ekpor, C. F., Okoli, I. C. and Iheukwumere, F. C. (2001): Performance and blood chemistry of Weaner Pigs Fed Raw *Mucuna* (Velvet Bean) meal. *Tropical.Animal Production Investigation* 4:49 – 54.
- Esonu, B. O., Udedibie, A. B. I. and Carlini, C. R. (1998).The effect of toasting, dry urea treatment and sprouting of some thermostable toxic factors in jack bean seed. *Nigerian Journal of Animal Production*, 25:36-39.
- FAO. (1992): Trade commerce of food and agric. organization of the United Nations, Rome, Vol.48.
- Goshit, D. T. (2015) Nutritive value of red kidney bean (*phaseolus vulgaris*) seed meal in broiler chickens diet, PhD. Thesis.
- Hang, D.T., and Preston, T.R. (2010). Effects of processing taro leaves on oxalate concentration as using the ensiled leaves as a protein source in pig diets in central Vietnam. *Livestock Research for Rural Development*, 22(68). Available: <http://www.rrd.org/lrrd22/4/hang22068.htm>

- Khare, S. K. (2000). Application of immobilized enzymes in soya bean, processing and utilization conference (ISPCRC 111): 2000 of the innovation era for soya bean 15-20, October, 2000. Isukuba, Ibaraka, Japan Pp.381-382.
- Krupa, U. (2008). Main Nutritional and anti-nutritional compounds of bean seeds: A review. *Polish Science*, 58(2): 149.
- Liener, I. E. (1980). Heat liable anti-nutritional factors. In: Advances in legume science. (Summer field, R. J. and Bunting, A. H. Eds.)New-London, Royal Botanic gardens. Pp. 157-170.
- Meyer, R. O., J. A. Froseth, and C. N. Coon (1992). Protein utilization and toxic effects of raw beans (*Phaseolus vulgaris*) for young pigs. *Journal of Animal Science* 55:1087-1098.
- Najime, D. (2003). Effect of processing on the utilization of Soya beans by broiler chickens. Unpublished M.Sc. Thesis Abubakar Tafawa Balewa University, Bauchi, Nigeria.
- Ologhobo, A. D., Apata, D. F. and Oyejide, A. (1993): Utilization of raw jackbean (*canavalia ensiformis*) and jackbean fractions in diets for broiler chicks. *British Poultry Science* 34, 323 – 337.
- Ologhobo, A. D., Mosenthin, R. and Alaka, O. O. (2003): Histological alterations in the internal organs of growing chicks from feeding raw jackbean or lima bean seeds. *Veterinary Human Toxicology*. 45(1): 10 – 13.
- Olomu, J. M. (1995). Feeding Management of poultry. In: monogastric animal nutrition principles and practices. A Jachen Publishers, Nigeria: 19-104; 108-118.
- Olomu, J. M. (2011). Monogastric animal nutrition (principles and practices). 2<sup>nd</sup> Edition St. Jackson Publishing Benin City Pp.206-228.
- Omage, J. J., Umar, I. A. and Bawa, G. S. (2002).Effect of Sesame (*Sesamum indicum L.*) Seed Oil on Blood and liver Lipid/Cholesterol Levels of Rats Fed a High Fat Diet. *Nigerian Journal of Experimental and Applied Biology*. 3:125 – 129.
- Pauzenga, U. (1985). Feeding Parent Stock. Zootech characteristics of weaning rabbits fed Graded levels. *International*, (34):22-25
- Reed, J, D (2001): Condensed Tannins, A factors limiting the using of cassava forage. *Journal of Science Food Agric* 33:213 – 217.
- Serres, J. L. (1999). Comparative nutritional evaluation of full fat soya-bean and source oil seeds. *Tropical Oil Seeds Journal*, 3(2): 100-105.
- Sonaiya, E. B. William, A. R. and S. A. Obi.(1986). A Biological and Economic Appraisal of Broiler Production up to 16 Weeks. *Journal of Animal Production Research*. 691: 73 – 79
- Steel, R. G. D. and Torrie, J. H. (1980). Principles and procedures of Statistics. A biometric Approach. 2<sup>nd</sup> edition McGraw-Hill Book Co., Inc. New York.
- Udedibie, A. B. I. and C. R. Carlini, (1997). Questions and answers to edibility problem of the carnivalia ensiformis seed. A review of Animal Feed Science and Technology, 74(2):95-106.
- Udedibie, A. B. I. and C. R. Carlini, (2000). Relative effects of dry and moist heat treatment of hemagglutinating and anti-tryptic activities of selected legume grains. *Nigerian Journal of Poultry Science*, 1:81-87.

British Journal of Multidisciplinary and Advanced Studies:

*Agriculture, 4(4),63-76, 2023*

Print ISSN: 2517-276X

Online ISSN: 2517-2778

Website: <https://bjmas.org/index.php/bjmas/index>

---

Published by European Centre for Research Training and Development-UK

Wu, W., Woodie, P. and Williams, M. (1996): Amino acid availability – corrected amino acid score of red kidney beans (*Phaseolus vulgaris L*). *Journal of Agriculture and Food Chemistry* 44:1296 – 1301.